


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




4126
4127
4128
4129
4130
4131
4132
4133
4134
4135
4136
4137
4138
4139
4140
4141
4142
4143
4144

Jem. Functional aspects of the photosynthetic light reactions in heat stressed Arabidopsis deficient in digalactosyl-diacylglycerol. <i>Journal of Plant Physiology</i> . 2011. 168:1526-1533
Philippe Frankart Eullaffroy. Energy fluxes and driving forces for photosynthesis in <i>Lemna minor</i> exposed to herbicides. <i>Aquatic Botany</i> . 2009. 90:172-178
Gary D. Smith Ewart. Immunochemical analysis of the soluble hydrogenase from the cyanobacterium <i>Anabaena cylindrica</i> . <i>Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology</i> . 1989. 997:83-89
Isabel Vitorica Fabregat. The pentose phosphate cycle is regulated by NADPH/NADP ratio in rat liver. <i>Archives of Biochemistry and Biophysics</i> . 1985. 236:110-118
A. Achak Farahi. Silver particles-modified carbon paste electrodes for differential pulse voltammetric determination of paraquat in ambient water samples. <i>Journal of the Association of Arab Universities for Basic and Applied Sciences</i> . 2016. 19:37-43
A. El Gaini Farahi. Interaction study of paraquat and silver electrode using electrochemical impedance spectroscopy: Application in milk and tomato samples. <i>Food Control</i> . 2015. 47:679-685
A. Lahrich Farahi. Parameters affecting the determination of paraquat at silver rotating electrodes using differential pulse voltammetry. <i>Analytical Chemistry Research</i> . 2014. <HT>1</HT>:16-21
J. A. Ebert Farrington. Bipyridylium quaternary salts and related compounds. V. Pulse radiolysis studies of the reaction of paraquat radical with oxygen. Implications for the mode of action of bipyridyl herbicides. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1973. 314:372-381
Dai-Jun Wang Feng. Hydrogen bonding-driven elastic bis(zinc)porphyrin receptors for neutral and cationic electron-deficient guests with a sandwich-styled complexing pattern. <i>Tetrahedron Letters</i> . 2007. 48:6181-6185
Victor M. Hatchikian Fernandez. Properties and reactivation of two different deactivated forms of <i>Desulfovibrio gigas</i> hydrogenase. <i>Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology</i> . 1985. 832:69-79
Y. Anglade Fernandez. Potentiation of iron-induced lipid peroxidation by a series of bipyridyls in relation to their ability to reduce iron. <i>Toxicology Letters</i> . 1997. 93:65-71
Roger J. Fido. Isolation and partial amino acid sequence of domains of nitrate reductase from spinach. <i>Phytochemistry</i> . 1991. 30:3519-3523
C. Eullaffroy Frankart. Comparative effects of four herbicides on non-photochemical fluorescence quenching in <i>Lemna minor</i> . <i>Environmental and Experimental Botany</i> . 2003. 49:159-168
Koyu Chan Fujita. Anaerobic purification, characterization and preliminary mechanistic study of recombinant nitrous oxide reductase from <i>Achromobacter cycloclastes</i> . <i>Journal of Inorganic Biochemistry</i> . 2007. 101:1836-1844
Carmine Caruso Gaeta. p-Sulfonatocalix[7]arene: synthesis, protolysis, and binding ability. <i>Tetrahedron</i> . 2008. 64:5370-5378
Lu. Decreasing the toxicity of paraquat through the complexation with sodium salicylate: Stoichiometric analysis. <i>Toxicology</i> . 2015. 336:96-98
D. W. Reif. Ferritin as a source of iron for oxidative damage. <i>Free Radic Biol Med</i> . 1992. 12:417-27
Harry W. Shen Gibson. Polymeric molecular shuttles: Polypseudorotaxanes & polyrotaxanes based on viologen (paraquat) urethane backbones & bis(p-phenylene)-34-crown-10. <i>Polymer</i> . 2014. 55:3202-3211
Ivan N. Gogotov. Relationships in hydrogen metabolism between hydrogenase and nitrogenase in phototrophic bacteria. <i>Biochimie</i> . 1978. 60:267-275

Plants are often submitted, in their natural environment, to various abiotic stresses such as heat stress. However, elevated
Analysis of fast chlorophyll fluorescence rise OJIP was carried out to assess the impact of diuron, paraquat and flazasulfuron
Antibodies were raised in mice against the 42 kDa subunit of the soluble hydrogenase purified from the cyanobacterium
The changes in the activity of the pentose phosphate cycle produced by the activation or inhibition of different NADPH-c
This paper describes the construction of silver particles-impregnated carbon paste electrode (Ag-CPE). The new electrode
This paper reports on the use of electrochemical impedance spectroscopy (EIS) for analytical determination of paraquat in
The electrochemical determination of aqueous paraquat PQ(II) by differential pulse voltammetry at a solid rotating silver
1. Rate constants for reduction of paraquat ion (1,1'-dimethyl-4,4'-bipyridyl-lum, PQ2+) to paraquat radical (PQ+
This Letter reports the design and synthesis of a new type of hydrogen bonding-mediated foldamer-derived tweezer rece
It has previously been shown that Desulfovibrio gigas hydrogenase, as isolated, has a relatively low activity in the hydrog
We determine the relative abilities of three bipyridyls (Paraquat P ++, Benzylviologen B++ and Diquat D++) to stimulate in
Fragments of spinach nitrate reductase (NR) were prepared by limited proteolysis of immunopurified enzyme using both
Aquatic ecosystems are exposed to an increasing contamination of pesticides such as herbicides through water runoff. Th
An overexpression system for nitrous oxide reductase (N2OR), an enzyme that catalyzes the conversion of N2O to N2 and
Water-soluble p-sulfonatocalix[7]arene 1 has been synthesized in good yield through standard procedures and its confor
Over the last decades, paraquat (1,1'-dimethyl-4,4'-bipyridilium dichloride; PQ) has been involved in numerous fatalities
The generation of deleterious activated oxygen species capable of damaging DNA, lipids, and proteins requires a catalyst
Reaction of di(p-isocyanatophenyl)methane (MDI, 4) with N,N'-di(2-hydroxyethyl)- (1b) or N,N'-di[2-(2'-hydroxyethoxy)]
Summary Purple bacteria Rhodospirillum rubrum and Thiocapsa roseopersicina form two enzymes, hydrogenase and nit

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



4145
4146
4147
4148
4149
4150
4151
4152
4153
4154
4155
4156
4157
4158
4159
4160
4161
4162
4163

Carlos Bes G. Structural requirements for the electron transfer between a flavoprotein and viologens. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1994. 1187:236-240
R. H. Brooke-Taylor Gordonsmith. Structural requirements of compounds to inhibit pulmonary diamine accumulation. <i>Biochemical Pharmacology</i> . 1983. 32:3701-3709
Zafirir Lapidot Goren. Photocatalysed regeneration of NAD(P)H by CdS and TiO ₂ semiconductors: applications in enzymatic synthesis. <i>Journal of Molecular Catalysis</i> . 1988. 47:21-32
van de Ven Govindjee. Chlorophyll a fluorescence lifetime distributions in open and closed Photosystem II reaction center preparations. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1990. 1015:173-179
Xu Govindjee. Chloroacetates as inhibitors of photosystem II: Effects on electron acceptor side. <i>Journal of Photochemistry and Photobiology B: Biology</i> . 1997. 37:107-117
E. Shavit Gross. Synthalin: An inhibitor of energy transfer in chloroplasts. <i>Archives of Biochemistry and Biophysics</i> . 1968. 127:224-228
Maxwell J. Johnston Gunter. A porphyrin-based crown ether co-receptor for the complexation of paraquat. <i>Tetrahedron Letters</i> . 1990. 31:4801-4804
Jessica Pandelia Hadj-Sa. The Carbon Monoxide Dehydrogenase from <i>Desulfovibrio vulgaris</i> . <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 2015. 1847:1574-1583
Achim Stransky Hager. Uncoupling of Photophosphorylation by Triethyl Lead (Et ₃ Pb ⁺) Generated from Tetraethyl Lead (Et ₄ Pb) in Illuminated Chloroplasts. <i>Journal of Plant Physiology</i> . 1989. 134:8-May
Marja Tuominen Hakala. Evidence for the role of the oxygen-evolving manganese complex in photoinhibition of Photosystem II. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 2005. 1706:68-80
D. O. Reeves Hall. Photosynthetic control in isolated spinach chloroplasts with endogenous and artificial electron acceptors. <i>Biochemical and Biophysical Research Communications</i> . 1971. 43:359-366
M. Helen Prince Hall. EPR spectroscopy of the iron-sulphur cluster and sirohaem in the dissimilatory sulphite reductase (desulphoviridin) from <i>Desulphovibrio gigas</i> . <i>Biochimica et Biophysica Acta (BBA) - Protein Structure</i> . 1979. 581:27-33
Randolph P. Roseboom Happe. Unusual FTIR and EPR properties of the H ₂ -activating site of the cytoplasmic NAD-reducing hydrogenase from <i>Ralstonia eutropha</i> . <i>FEBS Letters</i> . 2000. 466:259-263
Francis de Kouchkovsky Haraux. Further investigation on the lateral and transversal proton currents at the thylakoid membrane level by hydrogen-deuterium exchange. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1982. 679:235-247
Daniel C. Rinehart Harris. Reduction potential of iron in transferrin. <i>Biochimica et Biophysica Acta (BBA) - General Subjects</i> . 1985. 838:295-301
Jonathan J. Di Tomaso Hart. Characterization of the transport and cellular compartmentation of paraquat in roots of intact maize seedlings. <i>Pesticide Biochemistry and Physiology</i> . 1992. 43:212-222
H. Haseloff H. Free radical formation in chloroplasts. Methyl viologen action. <i>Journal of Photochemistry and Photobiology B: Biology</i> . 1992. 12:375-387
Maki Hasegawa Hayashi. Characterization of FMN-dependent NADH-quinone reductase induced by menadione in <i>Escherichia coli</i> . <i>Biochimica et Biophysica Acta (BBA) - General Subjects</i> . 1990. 1035:230-236
Takashi Hitomi Hayashi. New approach to the construction of an artificial hemoprotein complex. <i>Coordination Chemistry Reviews</i> . 1999. 190:961-974

The covalent binding of the viologen N-methyl-N ⁺ -(aminopropyl)-4,4'-bipyridinium (APMV) to the flavoprotein ferredoxin
The diamine, putrescine, is accumulated into slices of rat lung by a temperature and energy dependent process similar to
Photocatalysed regeneration of NAD(P)H is accomplished with CdS semiconductor powder and TiO ₂ colloids using forma
We have measured the decay of chlorophyll a fluorescence at 4
The two-electron gate of Photosystem II (PSII) is known to function by transferring electrons from the reduced one electr
Synthalin, decamethylene diguanidine, has been found to act as an energy-transfer inhibitor in chloroplasts. Both ATP for
A porphyrin strapped by a dibenzo- crown ether was synthesized and shown by ¹ H nmr spectroscopy to bind paraquat in
Ni-containing Carbon Monoxide Dehydrogenases (CODHs) catalyze the reversible conversion between CO and CO ₂ and a
Summary Tetraethyl lead (Et ₄ Pb), which is used as an anti-knock agent in gasoline, was transformed to the toxic triethyl l
Photoinhibition of PSII occurs at the same quantum efficiency from very low to very high light, which raises a question ab
Spinach chloroplasts, isolated rapidly in isotonic media will reproducibly give photosynthetic control rates (State 3/State
Desulphoviridin in the oxidized state showed EPR signals around g = 6, consistent with the sirohaem being in the high-spi
Soluble NAD-reducing [NiFe]-hydrogenase (SH) from <i>Ralstonia eutropha</i> (formerly <i>Alcaligenes eutrophus</i>) has an infrared
Energetically-coupled processes (electron flow, proton uptake and correlated pH gradient) were investigated on envelop
The reduction potential of Fe ³⁺ in transferrin was measured spectrophotometrically by equilibration with methyl violog
Uptake and compartmentation of paraquat was investigated in intact roots of hydroponically grown maize (<i>Zea mays</i> L.)
The generation of free radicals under various conditions in the presence of methyl viologen (MV ²⁺) was investigated in c
It was found that when <i>Escherichia coli</i> is grown in the presence of 0.2
Modified prosthetic metalloporphyrin, having a total of eight carboxylate groups at the terminal of two peripheral propic

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant